

# Galactic spiral patterns and dynamo action II: Asymptotic solutions

Luke Chamandy<sup>1</sup>, Kandaswamy Subramanian<sup>1</sup> & Anvar Shukurov<sup>2</sup> \*

<sup>1</sup>*Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411007, India*

<sup>2</sup>*School of Mathematics & Statistics, Newcastle University, Newcastle upon Tyne NE1 7RU*

Submitted to MNRAS

## ABSTRACT

The exploration of mean-field galactic dynamos affected by a galactic spiral pattern, begun in Chamandy et al. (2013, hereafter Paper I) with numerical simulations, is continued here with an asymptotic solution. The effect of the spiral pattern on the dynamo considered is the enhancement of the  $\alpha$  effect in spiral arms (which may overlap the gaseous spiral arms or be located in the interarm regions). The axisymmetric and enslaved non-axisymmetric modes of the mean magnetic field are studied semi-analytically to clarify and strengthen the numerical results. The mean-field dynamo model used generalizes the standard theory to include the delayed response of the mean electromotive force to variations of the mean magnetic field and turbulence parameters (the temporal non-locality, or  $\tau$  effect). Good qualitative agreement is obtained between the asymptotic solution presented and numerical solutions of Paper I for a global rigidly rotating material spiral (density wave). We find magnetic arms displaced from the  $\alpha$ -arms away from the co-rotation radius, so that the ridges of magnetic field strength are more tightly wound than the  $\alpha$ -arms. Moreover, the effect of a finite dynamo relaxation time  $\tau$  is to phase-shift the magnetic arms in the direction opposite to the galactic rotation. This mechanism can be used to explain the observed phase shifts between magnetic and material arms in some galaxies.

**Key words:** magnetic fields – MHD – galaxies: magnetic fields – galaxies: spiral – galaxies: structure – galaxies: ISM

## 1 INTRODUCTION

Disc galaxies typically contain regular (or large-scale or mean) magnetic fields of strength  $\sim 1 - 10 \mu\text{G}$ . In many cases, the strength of the non-axisymmetric components can be comparable to that of the axisymmetric component (Fletcher 2010). The non-axisymmetric part of the field takes the form of ‘magnetic spiral arms’, wherein the regular field is enhanced (Beck et al. 1996; Shukurov 2005; Beck 2012). Magnetic arms cannot be easily accounted for by mean-field dynamo theory if the underlying disc is assumed to be axisymmetric (Chamandy et al. 2013, hereafter Paper I). Moreover, they seem to be related (though in a non-trivial way) to the material (gaseous) spiral arms. Hence the need to develop a theory which relates the non-axisymmetry of the regular magnetic field to that of the underlying disc. This was the general aim of Paper I, where we approached the problem from a numerical standpoint and considered not only the linear growth of the field (kinematic

regime) but also the non-linear saturation phase. Here we take an analytical approach and therefore restrict the investigation to the kinematic regime. We are motivated, in part, by insights that were provided by analytical treatments in the past (Ruzmaikin et al. 1988, hereafter RSS88, for the axisymmetric case and Mestel & Subramanian 1991, hereafter MS91, for the non-axisymmetric case). The latter work was focused on how the  $|m| = 1$  (bisymmetric) azimuthal modes can be generated in a galactic disc with a two-arm spiral pattern. Numerical models (e.g. Moss 1998; Rohde et al. 1999; Paper I) show, however, that  $m = 0$  (axisymmetric) and  $|m| = 2$  (quadraxisymmetric) modes tend to be more prevalent in such discs (or  $m = 0$  and  $|m| = n$  for discs with  $n$  spiral arms). An analytical model that can explain such modes is therefore needed.

Our aim in this paper is thus a systematic interpretation of the key numerical results of Paper I (Chamandy et al. 2013) by solving a suitably simplified but essentially the same mean-field dynamo equations with an approximate semi-analytical method. Here we consider both axially symmetric and enslaved non-axisymmetric modes of a kinematic dynamo. The enslaved modes are those that have the same

\* E-mail: lchamandy@gmail.com (LC); kandu@iucaa.ernet.in (KS); anvar.shukurov@newcastle.ac.uk (AS)

growth rate as the leading axisymmetric mode; they occur because of deviations of the galactic disc from axial symmetry, e.g., due to a spiral pattern in the interstellar gas. For a two-armed spiral, the enslaved modes include the  $|m| = 2$  modes that corotate with the spiral pattern, as well as other, weaker, even- $m$  corotating modes. We leave the  $|m| = 1$  modes and other odd- $m$  corotating modes for a future work. Our model differs from earlier analytical study of MS91 in that here we:

(i) provide an asymptotic treatment of the  $m = 0$  and even- $m$  modes forced by a rigidly rotating two-arm spiral;

(ii) incorporate the minimal- $\tau$  approximation (MTA) and explore the effects of a finite dynamo relaxation time.

The plan of the paper is as follows. The basic equations and mathematical approach are summarized in Section 2. In Section 3, we present an asymptotic solution for the axisymmetric and enslaved non-axisymmetric magnetic modes which inhabit a disc containing a global, steady, rigidly rotating spiral density wave. We then compare our results with numerical solutions in Section 4. Conclusions and discussion for both Paper I and this paper can be found in Sections 5 and 6.

## 2 THE MEAN-FIELD DYNAMO AND THE TELEGRAPH EQUATION

The basic equation solved here is a generalization of the standard mean-field dynamo equation, and is derived in Paper I,

$$\tau \frac{\partial^2 \bar{\mathbf{B}}}{\partial t^2} + \frac{\partial \bar{\mathbf{B}}}{\partial t} = \tau \nabla \times \left( \bar{\mathbf{U}} \times \frac{\partial \bar{\mathbf{B}}}{\partial t} \right) + \nabla \times (\bar{\mathbf{U}} \times \bar{\mathbf{B}} + c_\tau \alpha \bar{\mathbf{B}}) + c_\tau \eta_t \nabla^2 \bar{\mathbf{B}}. \quad (1)$$

Here we use the same (standard) notation as in Paper I. We have neglected the terms  $\tau \eta \nabla^2 (\partial \bar{\mathbf{B}} / \partial t)$  and  $\eta \nabla^2 \bar{\mathbf{B}}$  on the right-hand side because of the high Ohmic conductivity of the interstellar gas,  $\eta \ll \eta_t$ . Following the usual mean-field approach, the velocity and magnetic fields,  $\mathbf{U}$  and  $\mathbf{B}$ , have each been written as the sum of an average part and a random part,

$$\mathbf{B} = \bar{\mathbf{B}} + \mathbf{b} \quad \text{and} \quad \mathbf{U} = \bar{\mathbf{U}} + \mathbf{u},$$

where overbar represents the ensemble average but for practical purposes spatial averaging over scales larger than the turbulent scale but smaller than the system size can be used (Germano 1992; Eyink 2012; Gent et al. 2012).

Equation (1) arises when the response time  $\tau$  of the mean electromotive force (emf)  $\mathcal{E} = \mathbf{u} \times \bar{\mathbf{b}}$  to changes in the mean field or small-scale turbulence is not negligible (e.g. Rheinhardt & Brandenburg 2012). The  $\tau$  effect can have various interesting astrophysical implications (see, e.g., Hubbard & Brandenburg 2009, which also contains a brief review of applications from the literature). A convenient approach to allow for this effect is the minimal- $\tau$  approximation (MTA) (Rogachevskii & Kleeorin 2000; Blackman & Field 2002; Brandenburg & Subramanian 2005a), which leads to a dy-

namical equation for  $\mathcal{E}$ :

$$\frac{\partial \mathcal{E}}{\partial t} = \frac{1}{\tau_c} (\alpha \bar{\mathbf{B}} - \eta_t \nabla \times \bar{\mathbf{B}}) - \frac{\mathcal{E}}{\tau}. \quad (2)$$

In the kinematic limit, and for isotropic and homogeneous turbulence, the turbulent transport coefficients are given by

$$\alpha = -\frac{1}{3} \tau_c \overline{\mathbf{u} \cdot \nabla \times \mathbf{u}}, \quad \eta_t = \frac{1}{3} \tau_c \overline{\mathbf{u}^2}, \quad (3)$$

with  $\tau_c$  the correlation time of the turbulence and  $\tau$  the relaxation time, found in numerical simulations to be close to  $\tau_c$  (Brandenburg & Subramanian 2005a,b, 2007). We retain  $c_\tau = \tau / \tau_c$  as a free parameter, but set it to unity in illustrative examples. For the sake of simplicity, we take  $\tau$  to be independent of scale and constant in space and time. Eq. (2), when combined with the mean-field induction equation, leads to Eq. (1).

The more standard mean-field dynamo equation

$$\frac{\partial \bar{\mathbf{B}}}{\partial t} = \nabla \times (\bar{\mathbf{U}} \times \bar{\mathbf{B}} + \alpha \bar{\mathbf{B}}) + \eta_t \nabla^2 \bar{\mathbf{B}}, \quad (4)$$

is recovered from Eq. (1) in the limit  $\tau \rightarrow 0$  with  $c_\tau = 1$ . Below, we refer to the approach leading to Eq. (4) as the ‘standard theory’.

## 3 ASYMPTOTIC SOLUTIONS FOR GALACTIC DYNAMOS

Many disc galaxies have a spiral structure which causes deviations from axial symmetry in both turbulent and regular gas flows, leading to non-axisymmetric magnetic fields. Here we follow MS91 by assuming that the  $\alpha$  effect is modulated by the spiral pattern. The nature of such a modulation is largely unexplored (Shukurov 1998; Shukurov & Sokoloff 1998), but it is reasonable to assume that  $\alpha$  is enhanced along a spiral perhaps overlapping with the gas spirals. The spiral patterns observed in the light of young stars are believed to represent a global density wave rotating at a fixed angular frequency, or possibly a superposition of such waves, with differing azimuthal symmetries and angular frequencies (e.g. Comparetta & Quillen 2012; Roškar et al. 2012). At least in some galaxies, they can be transient features (e.g. Dobbs et al. 2010; Sellwood 2011; Quillen et al. 2011; Wada et al. 2011; Grand et al. 2012) whose effect on the mean-field dynamo is discussed in Paper I. In this paper, we consider an enduring spiral with  $n$  arms and a constant pattern speed  $\Omega$ .

### 3.1 Basic equations

Since we are interested in non-axisymmetric magnetic fields that corotate with the spiral pattern, it is preferable to work in the corotating frame where the dynamo forcing is independent of time. Following MS91, we carry out a co-ordinate transformation from the inertial cylindrical frame  $\Sigma = (r, \phi, z, t)$  (with disc rotation axis as the  $z$ -axis) to the frame  $\tilde{\Sigma} = (\tilde{r}, \tilde{\phi}, \tilde{z}, \tilde{t})$  rotating with the pattern angular velocity  $\Omega$  (assumed constant in both time and position):

$$\begin{aligned} \tilde{\phi} &= \phi - \Omega t, & \tilde{r} &= r, & \tilde{z} &= z, & \tilde{t} &= t, \\ \tilde{\bar{\mathbf{B}}}_r &= \bar{\mathbf{B}}_r, & \tilde{\bar{\mathbf{B}}}_\phi &= \bar{\mathbf{B}}_\phi, & \tilde{\bar{\mathbf{B}}}_z &= \bar{\mathbf{B}}_z. \end{aligned} \quad (5)$$

so that

$$\left(\frac{\partial}{\partial\phi}\right)_t = \left(\frac{\partial}{\partial\tilde{\phi}}\right)_{\tilde{t}}, \quad \left(\frac{\partial}{\partial t}\right)_\phi = \left(\frac{\partial}{\partial\tilde{t}}\right)_{\tilde{\phi}} - \Omega \left(\frac{\partial}{\partial\tilde{\phi}}\right)_{\tilde{t}}. \quad (6)$$

The coordinate transformation shifts the mean velocity,  $\tilde{U}_\phi = \bar{U}_\phi - r\Omega$ , but the random velocity is left unchanged,  $\tilde{\mathbf{u}} = \mathbf{u}$ , as are both mean and random magnetic field. Therefore,  $\tilde{\alpha} = \alpha$  and  $\tilde{\eta}_t = \eta_t$ .

We use the thin-disc approximation,  $\partial/\partial z \gg \partial/\partial r$  and, eventually, consider tightly wound magnetic spirals,  $\partial/\partial r \gg \partial/r\partial\phi$ . Thus, we consider magnetic fields whose radial scale is asymptotically intermediate between the scale height of the galactic disc (of order 0.5 kpc) and its radial scale length (of order 10 kpc). We also adopt the  $\alpha\omega$ -dynamo approximation where the rate of production of the azimuthal magnetic field by the  $\alpha$  effect is negligible in comparison with the effect of the differential rotation. The mean velocity field is taken to be axisymmetric, purely azimuthal, and constant in time,

$$\bar{\mathbf{U}} = r\omega(r)\hat{\phi}, \quad (7)$$

where  $\hat{\phi}$  is the unit azimuthal vector. In the rotating frame, the gas angular velocity is  $\tilde{\omega}(r) = \omega(r) - \Omega$ . The magnitude of the rotational velocity shear is quantified with  $G(r) = r d\omega/dr = rd\tilde{\omega}/dr$ . Applying Eqs. (5) and (6) to the  $r$ - and  $\phi$ -components of Eq. (1), and dropping tildes for ease of notation (except for on  $\tilde{\omega}$ ), leads to the following equations in the corotating frame:

$$\begin{aligned} \hat{\mathcal{L}}\bar{B}_r = & -c_\tau \frac{\partial}{\partial z}(\alpha\bar{B}_\phi) \\ & + c_\tau \eta_t \left( \tilde{\nabla}^2 \bar{B}_r + \frac{1}{r^2} \frac{\partial^2 \bar{B}_r}{\partial \phi^2} - \frac{2}{r^2} \frac{\partial \bar{B}_\phi}{\partial \phi} \right), \end{aligned} \quad (8)$$

$$\begin{aligned} \hat{\mathcal{L}}\bar{B}_\phi = & \left( 1 + \tau \frac{\partial}{\partial t} - \Omega \tau \frac{\partial}{\partial \phi} \right) (G\bar{B}_r) \\ & + c_\tau \eta_t \left( \tilde{\nabla}^2 \bar{B}_\phi + \frac{1}{r^2} \frac{\partial^2 \bar{B}_\phi}{\partial \phi^2} + \frac{2}{r^2} \frac{\partial \bar{B}_r}{\partial \phi} \right), \end{aligned} \quad (9)$$

where

$$\hat{\mathcal{L}} = \left( 1 + \tau \frac{\partial}{\partial t} - \Omega \tau \frac{\partial}{\partial \phi} \right) \left( \frac{\partial}{\partial t} + \tilde{\omega} \frac{\partial}{\partial \phi} \right)$$

and

$$\tilde{\nabla}^2 X = \frac{\partial^2 X}{\partial z^2} + \frac{\partial}{\partial r} \left[ \frac{1}{r} \frac{\partial}{\partial r} (rX) \right].$$

These equations agree with Eqs. (2.2) and (2.3) of MS91 in the limit  $\tau \rightarrow 0$  with  $c_\tau = 1$ . Since, in the thin-disc approximation, both equations do not include  $\bar{B}_z$ , the equation for  $\bar{B}_z$  can be replaced by the solenoidality condition,

$$\frac{1}{r} \frac{\partial}{\partial r} (r\bar{B}_r) + \frac{1}{r} \frac{\partial \bar{B}_\phi}{\partial \phi} + \frac{\partial \bar{B}_z}{\partial z} = 0. \quad (10)$$

We emphasize that the governing equations acquire additional terms when transformed to the rotating frame whenever  $\tau$  is finite. On the contrary, the standard mean-field dynamo equation (4) is invariant under the transformation to a rotating frame. This can be understood physically in the following way. In the standard theory,  $\mathcal{E}$  at time  $t_0$  and position  $\mathbf{x}$  depends on  $\bar{\mathbf{B}}$  and the turbulence parameters at the same time and position. This is not true in the MTA, where  $\mathcal{E}(t_0, \mathbf{x})$  depends on the history of  $\bar{\mathbf{B}}(t, \mathbf{x})$ ,

$\tilde{\alpha}$  and  $\tilde{\eta}_t$  over, roughly,  $t_0 - \tau < t \leq t_0$ . [Here  $\tilde{\alpha}$  and  $\tilde{\eta}_t$  refer to the more general version of Eq. (2), wherein we have made the standard assumption  $\tilde{\alpha} \approx \alpha/\tau_c$ ,  $\tilde{\eta}_t \approx \eta_t/\tau_c$  (Brandenburg & Subramanian 2005a).] Thus,  $\mathcal{E}(t_0, \mathbf{x})$  is affected by how fast variations in the field and  $(\alpha, \eta_t)$  are sweeping past the position  $\mathbf{x}$ . In other words, the dynamo equations are no longer invariant under the transformation to a rotating frame.

### 3.2 Approximate solution

When the coefficients of Eqs. (8) and (9) depend on the azimuthal angle  $\phi$  as  $\exp(im\phi)$  with certain  $m$ , their solutions can be represented in the form

$$\begin{pmatrix} \bar{B}_r \\ \bar{B}_\phi \end{pmatrix} = \sum_{m=-\infty}^{\infty} \begin{pmatrix} a_m(r, z) \\ b_m(r, z) \end{pmatrix} \exp(im\phi + \Gamma t), \quad (11)$$

so that Eqs. (8) and (9) reduce to

$$\begin{aligned} \sum_{m=-\infty}^{\infty} e^{im\phi} \left[ \mathcal{L}a_m + c_\tau \frac{\partial}{\partial z}(\alpha b_m) \right. \\ \left. - c_\tau \eta_t \left( \tilde{\nabla}^2 a_m - m^2 \frac{a_m}{r^2} - 2im \frac{b_m}{r^2} \right) \right] = 0, \end{aligned} \quad (12)$$

$$\begin{aligned} \mathcal{L}b_m - (1 + \Gamma\tau - im\Omega\tau)Ga_m \\ - c_\tau \eta_t \left( \tilde{\nabla}^2 b_m - m^2 \frac{b_m}{r^2} + 2im \frac{a_m}{r^2} \right) = 0, \end{aligned} \quad (13)$$

where

$$\mathcal{L} = (1 + \Gamma\tau - im\Omega\tau)(\Gamma + im\tilde{\omega}).$$

We note that  $\alpha$  is, in general, a function of  $r$ ,  $\phi$  and  $z$ , whereas  $\omega$  and  $G$  are assumed to be functions of  $r$  alone. When obtaining Eq. (13) from Eq. (9), the summation can be dropped because the only  $\phi$ -dependence that occurs in this equation is in the common factor  $\exp(im\phi)$ , whereas  $\alpha$  depends on  $\phi$  in Eq. (12).

To make further progress, we model  $\alpha$  as an  $n$ -armed, rigidly rotating global spiral (thus, in the rotating frame it is time-independent),

$$\alpha = \alpha_0 [1 + \epsilon_\alpha \cos(n\phi - \kappa r)] \equiv \alpha_0 + \alpha_n e^{in\phi} + \alpha_{-n} e^{-in\phi}, \quad (14)$$

where

$$\alpha_n = \frac{1}{2}\alpha_0 \epsilon_\alpha e^{-i\kappa r}, \quad \alpha_{-n} = \frac{1}{2}\alpha_0 \epsilon_\alpha e^{i\kappa r} = \alpha_n^*, \quad (15)$$

where  $\alpha_0$  may depend on  $r$  and  $z$ , and asterisk denotes complex conjugate. With this convention,  $\kappa < 0$  describes a trailing Archimedean spiral. We choose an Archimedean rather than a logarithmic spiral in order to simplify the calculations; this choice does not affect our conclusions. The requirement that each coefficient of Eq. (12) vanishes gives

$$\begin{aligned} \mathcal{L}a_m - c_\tau \eta_t \left( \tilde{\nabla}^2 a_m - m^2 \frac{a_m}{r^2} - 2im \frac{b_m}{r^2} \right) \\ = -c_\tau \frac{\partial}{\partial z} (\alpha_0 b_m + \alpha_n b_{m-n} + \alpha_{-n} b_{m+n}). \end{aligned}$$

This implies that modes with  $m = \dots, -2n, -n, 0, n, 2n, \dots$  are coupled to each other, and likewise modes with  $m = \dots, -2n+1, -n+1, 1, n+1, 2n+1, \dots$ , etc. However, these sets of modes are decoupled from one another. Modes that are coupled to the generally dominant  $m = 0$  mode grow

along with it, and are thus called ‘enslaved’ modes. One naturally expects the lowest-order enslaved modes of order  $n$  to dominate the higher-order enslaved modes, as the former are forced directly. This was in fact borne out in the numerical work of Paper I. Therefore, the  $m = \pm n$  modes will be the dominant non-axisymmetric modes for a galaxy with an  $n$ -armed spiral pattern.

Taking  $n = 2$  (a two-armed  $\alpha$ -spiral), MS91 and Subramanian & Mestel (1993, hereafter SM93) truncated the series of coupled equations by neglecting  $|m| \geq 3$ . Thus, only the  $m = \pm 1$  are coupled and likewise  $m = 0$  is coupled to  $m = \pm 2$ . Here we truncate the series in the same way; the numerical solutions of Paper I confirm that this is an excellent approximation for much of the parameter space (however, see below). The case of  $m = \pm 1$  has already been considered in MS91. Thus, we focus on the  $m = 0, \pm 2$  case here, for which the governing equations are

$$(1 + \Gamma\tau)\Gamma a_0 - c_\tau \eta_t \tilde{\nabla}^2 a_0 = -c_\tau \frac{\partial}{\partial z}(\alpha_0 b_0 + \alpha_2 b_{-2} + \alpha_{-2} b_2), \quad (16)$$

$$(1 + \Gamma\tau)\Gamma b_0 - c_\tau \eta_t \tilde{\nabla}^2 b_0 = (1 + \Gamma\tau)Ga_0, \quad (17)$$

while, for  $m = \pm 2$  and  $n = 2$ ,

$$\begin{aligned} La_{\pm 2} - c_\tau \eta_t \left( \tilde{\nabla}^2 a_{\pm 2} - 4 \frac{a_{\pm 2}}{r^2} \mp 4i \frac{b_{\pm 2}}{r^2} \right) \\ = -c_\tau \frac{\partial}{\partial z}(\alpha_0 b_{\pm 2} + \alpha_{\pm 2} b_0), \end{aligned} \quad (18)$$

$$\begin{aligned} Lb_{\pm 2} - c_\tau \eta_t \left( \tilde{\nabla}^2 b_{\pm 2} - 4 \frac{b_{\pm 2}}{r^2} \pm 4i \frac{a_{\pm 2}}{r^2} \right) \\ = (1 + \Gamma\tau \mp 2i\Omega\tau)Ga_{\pm 2}, \end{aligned} \quad (19)$$

where

$$L = (1 + \Gamma\tau \mp 2i\Omega\tau)(\Gamma \pm 2i\tilde{\omega}).$$

### 3.2.1 Non-axisymmetric modes under the no- $z$ and WKBJ approximations

We note that for the even modes  $m = 0, \pm 2$  there are six coupled partial differential equations. MS91 solved the corresponding simpler problem for the odd ( $m = \pm 1$ ) modes, which involves four coupled PDEs, using WKBJ methods, but they did not do the same for the even modes. Thus we would like to get analytical insights into these perhaps more important modes. Therefore we proceed as follows: Our plan is to first solve for  $a_{\pm 2}$  and  $b_{\pm 2}$ . We do this by using the no- $z$  approximation for the  $z$ -derivatives (SM93; Moss 1995; Phillips 2001), and a WKBJ-type approximation to handle radial diffusion. This yields algebraic equations which can be solved to obtain explicit expressions for  $a_{\pm 2}$  and  $b_{\pm 2}$ . We then substitute them into the  $m = 0$  equations, (16) and (17), and apply the no- $z$  approximation to obtain Schrödinger-type differential equations in  $r$  with the effective potential  $V(r)$ . Bound states in this potential correspond to exponentially growing solutions, and these are obtained using the WKBJ approximation. This is done iteratively using numerical methods, so that the method is semi-analytical rather than analytical. Thus we avoid solving the six coupled differential equations (16)–(19). We shall see that this procedure, in which four of the six differential

equations are approximated as algebraic equations, works well for the case where the vertical turbulent diffusion is much stronger than the radial turbulent diffusion, i.e., for solutions of a large radial scale (wavelength) in a thin disc.

In the no- $z$  approximation, the vertical diffusion terms are approximated as

$$\frac{\partial^2 a_{\pm 2}}{\partial z^2} \simeq -\frac{\pi^2 a_{\pm 2}}{4h^2}, \quad \frac{\partial^2 b_{\pm 2}}{\partial z^2} \simeq -\frac{\pi^2 b_{\pm 2}}{4h^2},$$

where  $h$  is the disc half-thickness. Further, the terms containing  $\alpha$  can be approximated as

$$\begin{aligned} \frac{\partial}{\partial z}(\alpha_0 b_0 + \alpha_2 b_{-2} + \alpha_{-2} b_2) &\simeq \frac{2}{\pi h}(\alpha_0 b_0 + \alpha_2 b_{-2} + \alpha_{-2} b_2), \\ \frac{\partial}{\partial z}(\alpha_0 b_{\pm 2} + \alpha_{\pm 2} b_0) &\simeq \frac{2}{\pi h}(\alpha_0 b_{\pm 2} + \alpha_{\pm 2} b_0). \end{aligned}$$

For the radial diffusion terms, we start by representing  $a_{\pm 2}$  and  $b_{\pm 2}$  as

$$a_{\pm 2} = -\bar{a}(r)e^{\mp i\theta_a(r)}, \quad b_{\pm 2} = \bar{b}(r)e^{\mp i\theta_b(r)}, \quad (20)$$

where the minus sign in front of  $\bar{a}$  is introduced for future convenience, and where we have removed the  $z$ -dependence of  $\bar{a}$  and  $\bar{b}$  due to the fact that we are now working in the no- $z$  approximation. Note that  $|a_2| = |a_{-2}|$  and  $|b_2| = |b_{-2}|$ , whereas the phases of  $a_2$  and  $a_{-2}$  (or  $b_2$  and  $b_{-2}$ ) must be equal in magnitude and opposite in sign, since  $B_r$  and  $B_\phi$  are real. We then apply the WKBJ-type tight-winding approximation, assuming that the amplitude varies much less rapidly with radius than the phase, i.e.,

$$\left| \frac{d\bar{a}}{dr} \right| \ll \left| \bar{a} \frac{d\theta_a}{dr} \right|, \quad \left| \frac{d\bar{b}}{dr} \right| \ll \left| \bar{b} \frac{d\theta_b}{dr} \right|.$$

The radial diffusion terms are then approximated by

$$\frac{d}{dr} \left[ \frac{1}{r} \frac{d}{dr} (ra_{\pm 2}) \right] \simeq -(\theta_a'^2 \pm i\theta_a'') a_{\pm 2}, \quad (21)$$

$$\frac{d}{dr} \left[ \frac{1}{r} \frac{d}{dr} (rb_{\pm 2}) \right] \simeq -(\theta_b'^2 \pm i\theta_b'') b_{\pm 2}, \quad (22)$$

where prime stands for  $d/dr$ . It should be noted that in the region where the relative strength of non-axisymmetric modes with respect to axisymmetric modes is expected to be the largest (near the co-rotation radius), we expect  $1/r \ll |d/dr|$  since the co-rotation radius is typically comparable to the size of the galaxy. Thus, we have dropped terms proportional to  $1/r$ . Since turbulent diffusion terms containing the  $\phi$ -derivatives are proportional to  $1/r^2$ , they, too, can be neglected.

As shown in Appendix A, solutions of Eqs. (18) and (19) can be represented as

$$a_{\pm 2} = -\frac{|D_0|b_0\epsilon_\alpha}{2} \sqrt{\frac{\tilde{A}^2 + \tilde{B}^2}{A^2 + B^2}} \exp[\mp i(\kappa r + \beta + \tilde{\beta})], \quad (23)$$

$$b_{\pm 2} = \frac{|D_0|b_0\epsilon_\alpha}{2} \frac{1}{\sqrt{A^2 + B^2}} \exp[\mp i(\kappa r + \beta)], \quad (24)$$

where  $A$ ,  $B$ ,  $\tilde{A}$  and  $\tilde{B}$ , as well as  $\beta$  and  $\tilde{\beta}$ , are functions of position defined in Appendix A. Here the dynamo number is defined as

$$D_0 = \frac{\alpha_0 G h^3}{\eta_t^2} = \frac{\alpha_0 G t_d^2}{h} < 0, \quad (25)$$

where  $t_d = h^2/\eta_t$  is the vertical turbulent diffusion time scale.

Using Eqs. (20), we now make the identifications

$$\theta_a = \kappa r + \beta + \tilde{\beta}, \quad \theta_b = \kappa r + \beta,$$

and so

$$\theta'_a = \kappa + \beta' + \tilde{\beta}', \quad \theta'_b = \kappa + \beta', \quad \theta''_a = \beta'' + \tilde{\beta}'', \quad \theta''_b = \beta''. \quad (26)$$

Clearly,  $\beta$  and  $\tilde{\beta}$  depend on  $A$ ,  $B$ ,  $\tilde{A}$  and  $\tilde{B}$  (Eqs. A5 and A8), which, in turn, depend on the derivatives of  $\beta$  and  $\tilde{\beta}$  through  $\theta'_a$ ,  $\theta'_b$ ,  $\theta''_a$  and  $\theta''_b$  (Eqs. A6, A7, A9 and A10). This means that, unless  $\beta'$  and  $\tilde{\beta}'$  are completely neglected, it is not possible to obtain an analytical solution. We will see however that, if the terms involving  $\beta'$ ,  $\tilde{\beta}'$ ,  $\beta''$  and  $\tilde{\beta}''$  are small compared with the turbulent diffusion terms that do not vary with  $r$ , then  $\beta$  and  $\tilde{\beta}$  can be determined semi-analytically, by successive iteration.

The need for the iterations can be understood as follows. Firstly, if  $\beta'$  and  $\tilde{\beta}'$  were zero, then the radial phase would be  $\kappa r$ , which is the same as the radial phase of the  $\alpha$ -spiral. Thus, the magnetic arms would be as tightly wound as the  $\alpha$ -arms. In fact, the extra contributions to the radial phase  $\beta + \tilde{\beta}$  for  $\overline{B}_r$  and  $\beta$  for  $\overline{B}_\phi$ , are finite, and turn out to have the same sign as  $\kappa$ . This implies that the magnetic arms (defined as the set of positions where  $\overline{B}$  is maximum), are more tightly wound than the material arms that produce them. This is because the enhancement of field due to the enhanced dynamo action within the  $\alpha$ -arm is advected downstream with the differentially rotating gas. Ultimately a balance is set up between the competing effects of preferential magnetic field generation in the  $\alpha$ -arms and advection downstream, resulting in stationary magnetic arms that are more tightly wound than the  $\alpha$ -arms, and cross them near the co-rotation circle. Moreover, this balance, which determines the pitch angle of the magnetic arms (ridges), and thus the values of  $\beta$  and  $\tilde{\beta}$ , involves many quantities, including the derivatives of  $\beta$  and  $\tilde{\beta}$  through the radial diffusion (as seen in the equations of Appendix A). This self-dependence results from the fact that the radial diffusion is stronger when the magnetic arms (ridges) are more tightly wound, as can be seen from Eqs. (21) and (22). Thus, the radial diffusion feeds back onto itself, forcing us to iterate to obtain the functions  $\beta$  and  $\tilde{\beta}$ .

### 3.2.2 The global solution

Once  $a_{\pm 2}$  and  $b_{\pm 2}$  are obtained, we can solve Eqs. (16) and (17) for  $a_0$  and  $b_0$ . Following RSS88 and MS91 by invoking the thinness of the disc, we factorise the solution into the local ( $\tilde{a}$  and  $\tilde{b}$ ) and global ( $q$ ) parts,

$$a_0(r) = \tilde{a}q(r), \quad b_0(r) = \tilde{b}q(r). \quad (27)$$

Generally, the local solution depends on  $z$ , i.e.,  $\tilde{a}(z)$  and  $\tilde{b}(z)$ . However, the  $z$ -dependence has been removed using the no- $z$  approximation, and the local solution is a vectorial constant. Substituting Eqs. (23) and (24) into Eq. (11), and

using Eqs. (27), we can write the overall solution as

$$\overline{B}_r = \tilde{a}q \left[ 1 - \frac{\tilde{b}}{\tilde{a}} \sqrt{\frac{\tilde{A}^2 + \tilde{B}^2}{A^2 + B^2}} \epsilon_\alpha |D_0| \cos(2\phi - \kappa r - \beta - \tilde{\beta}) \right] e^{\Gamma t}, \quad (28)$$

$$\overline{B}_\phi = \tilde{b}q \left[ 1 + \frac{1}{\sqrt{A^2 + B^2}} \epsilon_\alpha |D_0| \cos(2\phi - \kappa r - \beta) \right] e^{\Gamma t}. \quad (29)$$

Magnetic field lines of the mean field are expected to be trailing spirals since  $G < 0$ , which implies  $\tilde{b}/\tilde{a} < 0$ . Therefore, the overall phase difference between  $\overline{B}_r$  and  $\overline{B}_\phi$  has magnitude  $\tilde{\beta}$  [without the minus sign introduced in front of  $\tilde{a}$  in Eq. (20), there would have been an extra phase difference  $\pi$ ]. The pitch angle of the magnetic field, defined as

$$p_B = \arctan \frac{\overline{B}_r}{\overline{B}_\phi}, \quad (30)$$

is independent of  $q$ .

Each component of the mean magnetic field consists of an axisymmetric part and that having the  $m = 2$  symmetry,

$$\overline{B}_i = \overline{B}_i^{(0)} + \overline{B}_i^{(2)} \cos[2(\phi - \phi_i)],$$

where  $i = r, \phi$ . The magnitude of the  $r$ -component of the field is maximum where  $2\phi - \kappa r - \beta - \tilde{\beta} = 0$ , while the magnitude of the  $\phi$ -component is maximum for  $2\phi - \kappa r - \beta = 0$ . On the other hand, the magnitude of  $\alpha$  is maximum where  $2\phi - \kappa r = 0$ . Therefore, the ( $\pi$ -fold degenerate) phase differences between the  $r$  and  $\phi$  components of the magnetic field and the  $\alpha$ -spiral are given, respectively, by

$$\Delta_r = \frac{\beta + \tilde{\beta}}{2}, \quad \Delta_\phi = \frac{\beta}{2}.$$

### 3.2.3 Growth rates of non-axisymmetric magnetic modes

We are now prepared to obtain the growth rate  $\Gamma$ . For this purpose we substitute Eq. (27) into the no- $z$  versions of Eqs. (16) and (17), and, in Eq. (16), we also substitute Eq. (A1) for  $b_{\pm 2}$ . In addition, we make use of the relation

$$\alpha_2 \alpha_{-2} = \alpha_0^2 \epsilon_\alpha^2 / 4,$$

obtained from Eq. (15). Equations (16) and (17) then reduce to

$$(1 + \Gamma\tau)\Gamma - c_\tau \frac{\eta_t}{q} \left[ \frac{(rq)'}{r} \right]' = -\frac{2c_\tau}{\pi h} \alpha_0 \frac{\tilde{b}}{\tilde{a}} \left[ 1 + \frac{|D_0| \epsilon_\alpha^2}{4} (X_2 + X_{-2}) \right] - c_\tau \eta_t \frac{\pi^2}{4h^2},$$

$$(1 + \Gamma\tau)\Gamma - c_\tau \frac{\eta_t}{q} \left[ \frac{(rq)'}{r} \right]' = (1 + \Gamma\tau)G \frac{\tilde{a}}{\tilde{b}} - c_\tau \eta_t \frac{\pi^2}{4h^2},$$

where  $X_{\pm 2}$  are defined in Eq. (A2). Note that  $\epsilon_\alpha = 0$  corresponds to an axisymmetric forcing of the dynamo. It is convenient to introduce a new quantity  $\gamma(r)$  defined as

$$\gamma(r) = (1 + \Gamma\tau)\Gamma - c_\tau \frac{\eta_t}{q} \left[ \frac{(rq)'}{r} \right]',$$

whose significance will shortly become evident. Now the equations can be separated into a global eigen-equation for  $q(r)$ , which only contains derivatives with respect to  $r$ , and two local equations for  $\tilde{a}$  and  $\tilde{b}$ , which generally only have

derivatives with respect to  $z$  but become algebraic equations under the no- $z$  approximation.

The global equation is given by

$$(1 + \Gamma\tau)\Gamma q = \gamma q + c_\tau \eta_t \left[ \frac{(rq)'}{r} \right]', \quad (31)$$

while the local equations can be rewritten as

$$\left( \gamma + \frac{\pi^2 c_\tau}{4t_d} \right) \tilde{a} = -c_\tau \frac{2\alpha_0}{\pi h} \tilde{b} \left[ 1 + \frac{|D_0| \epsilon_\alpha^2}{4} (X_2 + X_{-2}) \right], \quad (32)$$

$$\left( \gamma + \frac{\pi^2 c_\tau}{4t_d} \right) \tilde{b} = (1 + \Gamma\tau) G \tilde{a}. \quad (33)$$

It is now clear that  $\gamma$  is the local (at a given  $r$ ) growth rate of the dynamo modes in the slab approximation.

These homogeneous equations in  $\tilde{a}$  and  $\tilde{b}$  are easily solved to yield  $\tilde{b}/\tilde{a}$  and  $\gamma$ . From Eq. (33), we obtain

$$\frac{\tilde{b}}{\tilde{a}} = \frac{(1 + \Gamma\tau) G t_d}{\gamma t_d + \frac{1}{4} c_\tau \pi^2}.$$

Then the vanishing of the determinant of the coefficients of Eqs. (32) and (33) yields

$$\gamma = t_d^{-1} \left\{ -\frac{\pi^2 c_\tau}{4} \pm \sqrt{\frac{2c_\tau}{\pi} (1 + \Gamma\tau) |D_0| \left[ 1 + \frac{\epsilon_\alpha^2 |D_0| A}{2(A^2 + B^2)} \right]} \right\}, \quad (34)$$

where the positive sign in front of the square root provides growing solutions,  $\gamma > 0$ .

With  $\epsilon_\alpha = 0$  (axisymmetric forcing) and  $\gamma = 0$  (neutral stability of the mean magnetic field), we obtain an estimate for the local critical dynamo number, i.e., the minimum magnitude of the dynamo number for the field to grow rather than decay,

$$|D_{0,\text{cr}}| = \frac{\pi^5 c_\tau}{32(1 + \Gamma\tau)} \approx \frac{9.6 c_\tau}{1 + \Gamma\tau}.$$

It is slightly larger than the  $D_{0,\text{cr}} = 8$ , obtained by the numerical solution of the  $z$ -dependent versions of Eqs. (32) and (33) for  $c_\tau = 1$ ,  $\tau = \epsilon_\alpha = 0$  (RSS88); we note that  $\Gamma\tau \ll 1$ .

For  $\epsilon_\alpha = 0$ , the magnetic pitch angle, given by Eq. (30), can be expressed as

$$\tan p_{B,0} = \frac{\tilde{a}}{\tilde{b}} = -\sqrt{\frac{2c_\tau}{\pi(1 + \Gamma\tau)h} \frac{\alpha_0}{|G|}}. \quad (35)$$

A similar result can be obtained from perturbation theory (Sur et al. 2007). With F. Krause's law,

$$\alpha_0 = l^2 \omega / h, \quad (36)$$

and  $G = -\omega$  (flat rotation curve), this becomes

$$\tan p_{B,0} = -\frac{l}{h} \sqrt{\frac{2c_\tau}{\pi(1 + \Gamma\tau)}}.$$

The azimuthal average of the pitch angle in the kinematic regime obtained numerically in Paper I, where Eq. (36) was used, agrees very well with this formula. This can be seen by using the relevant parameters from that paper in the above equation, and then comparing with Fig. 8d of that paper.

However, here we shall take  $\alpha_0$  and  $h$  constant with radius, so that we have

$$\tan p_{B,0} = -\sqrt{\frac{2c_\tau}{\pi(1 + \Gamma\tau)} \frac{\alpha_0}{h\omega}}, \quad (37)$$

so that  $|p_{B,0}|$  now increases with radius, as  $\omega$  is a decreasing function of  $r$ .

Now, substituting Eq. (34) into Eq. (31), we obtain a Schrödinger-type equation for  $q(r)$ ,

$$-\nabla_r^2 q + V(r)q(r) = E q(r), \quad (38)$$

where  $\nabla_r^2 = (rq')'/r$  is the radial part of the Laplacian in cylindrical coordinates, with prime denoting  $\partial/\partial r$ . Here, the 'potential' is given by

$$V = \frac{1}{r^2} - \frac{\gamma t_d}{c_\tau h^2} = \frac{1}{r^2} + \frac{\pi^2}{4h^2} - \frac{1}{h^2} \sqrt{\frac{2}{\pi c_\tau} (1 + \Gamma\tau) |D_0| \left[ 1 + \frac{\epsilon_\alpha^2 |D_0| A}{2(A^2 + B^2)} \right]}, \quad (39)$$

and the 'energy' eigenvalue, by

$$E = -\frac{(1 + \Gamma\tau)\Gamma t_d}{c_\tau h^2}. \quad (40)$$

Note that bound states in the 'potential'  $V(r)$  with  $E < 0$  correspond to growing modes with  $\Gamma > 0$ . We solve this equation using the WKBJ theory: in terms of a scaled variable  $x$ , introduced via  $r = h e^x$  (with  $h = \text{const}$ ,  $-\infty < x < \infty$  and  $dx = dr/r$ ), Eq. (38) reduces to

$$\frac{d^2 q}{dx^2} + p(x)q(x) = 0,$$

where

$$p(x) = h^2 e^{2x} [E - V(x)].$$

The boundary conditions are  $q = 0$  at  $r = 0$  and  $r \rightarrow \infty$  (or  $x \rightarrow \pm\infty$ ). Suppose that  $p(x) = 0$  for  $x = x_\pm$  (corresponding to  $r_\pm$ ), with  $x_- < x_+$  (for our purposes, it is sufficient to consider the case of just two roots). The standard WKBJ theory yields the quantization condition

$$\int_{r_-}^{r_+} \frac{\sqrt{p(r)}}{r} dr = \int_{r_-}^{r_+} [E - V(r)]^{1/2} dr = \frac{\pi}{2} (2k + 1), \quad (41)$$

with  $k = 0, 1, 2, \dots$ , which can be used to obtain the growth rate  $\Gamma$ . Note, however, that  $\Gamma$  enters both the 'potential' and the 'energy', so the WKBJ approach has to be supplemented with an iteration procedure to converge on the correct value of  $\Gamma$ , in addition to the iterations over  $\beta$  and  $\tilde{\beta}$  discussed above.

As we will see below, the  $\phi$ -dependent term in the  $\alpha$  effect leads to a deepening of the potential well near the co-rotation radius, allowing non-axisymmetric modes that corotate with the  $\alpha$ -spiral to exist there. Outside this region (i.e., far from the co-rotation radius) axisymmetric modes will dominate, while non-axisymmetric modes will be weaker. Non-corotating non-axisymmetric modes (which rotate at approximately the local angular velocity at the radius where their eigenfunctions are maximum) are also expected to exist, but to be sub-dominant everywhere (MS91). The asymptotic solutions developed here apply to the corotating magnetic modes in the kinematic regime, but the numerical simulations of Paper I are, of course, not restrictive.




**Figure 1.** (a) The potential (39) of (38) for  $\tau = 2l/u$  (thick, black, solid curve) and  $\tau \rightarrow 0$  (thin, red, solid curve), near the co-rotation radius (dashed vertical line). The ‘energy’ eigenvalues (40) of (38) are represented by horizontal lines inside the wells. The axisymmetric part of the potential is also shown as a dashed line in both cases. (b) The solid curves represent the ratio of non-axisymmetric to axisymmetric parts of  $\overline{B}_\phi$  at the azimuth  $\phi = \phi_{\text{cor}} = \kappa r_{\text{cor}}/2$ , while the dashed curves show the corresponding magnitudes of the ratios  $\pm \overline{B}_\phi^{(2)}/\overline{B}_\phi^{(0)}$ . The dash-dotted curve shows the variation of  $\alpha(r, \phi_{\text{cor}})$ . (c) The phase differences,  $\Delta_\phi$  (solid) and  $\Delta_r$  (dashed), between the components of the magnetic spiral arms and the  $\alpha$ -spiral arms. The radius  $r_{\text{max}}$  where the envelope of  $\overline{B}_\phi^{(2)}/\overline{B}_\phi^{(0)}$  is maximum is shown as a dotted vertical line of the appropriate colour and thickness. (d) Pitch angle  $p_B$ , shown with the opposite sign, at  $\phi = \phi_{\text{cor}} = \kappa r_{\text{cor}}/2$  (solid). Also shown is  $-p_B$  for the purely axisymmetric case (dashed).

### 3.3 Illustrative example

We determine  $\Gamma$  and the properties of the mean magnetic field by varying  $\Gamma$  iteratively until the quantization condition (41) is satisfied for  $k = 0$  (the fastest growing mode). In addition, there is the complication that  $\beta$ ,  $\tilde{\beta}$ , and hence  $\theta_a$ ,  $\theta_b$ ,  $A$ ,  $B$ ,  $\tilde{A}$ ,  $\tilde{B}$ , etc., are not a priori known (see Section 3.2.1 and Appendix A). As mentioned above, we resolve this by iterating these variables as well. That is, we start with  $\beta'(r)$ ,  $\tilde{\beta}'(r)$  and their derivatives equal to zero, and at each iteration, we obtain these functions from Eqs. (A5) and (A8) to use them as input for the next iteration. The iterations converge if the terms involving  $\beta'$  and  $\tilde{\beta}'$  are small compared with the turbulent diffusion terms that do not vary with  $r$ . In the example below,  $\beta'$  and  $\tilde{\beta}'$  are comparable to  $\kappa$  near the co-rotation. However, it can be seen from Eqs. (A6), (A7), (A9) and (A10) that the terms involving the radial derivatives are not very important if the radial diffusion is much weaker than the vertical diffusion, that is if  $\theta_a'^2, \theta_b'^2 \ll \pi^2/4h^2$ , which may permit the iterations to converge. Therefore, in the example below, we choose the disc half-thickness  $h$  to be small enough (somewhat smaller than the standard value) that the iterations do in fact converge.

In order to determine the potential and growth rates of the corotating non-axisymmetric modes, we must first specify the functional forms of the scale height  $h(r)$ , angular velocity  $\omega(r)$  and  $\alpha_0(r)$ . For simplicity, we choose

$h = 0.1 \text{ kpc} = \text{const}$ . We also take  $\eta_t = ul/3$ , where  $u$  and  $l$  are the rms turbulent velocity and scale, and adopt  $l = h/4 = 0.025 \text{ kpc}$ , and  $u = 12ht_d^{-1} = 5 \text{ km s}^{-1}$ . These values are different from those usually adopted,  $u = 10 \text{ km s}^{-1}$  and  $l = 0.1 \text{ kpc}$ . Our choice is motivated by the desire to improve the convergence of the iterations since our aim here is mainly to illustrate the solution procedure and to discuss the qualitative properties of the eigen-solutions. However, we take care to keep the key parameter, the vertical turbulent diffusion time, at  $t_d = 0.24 \text{ Gyr}$ , similar to (about twice smaller than) the standard value.

As in Paper I, we use Brandt’s rotation curve,

$$\omega = \frac{\omega_0}{[1 + (r/r_\omega)^2]^{1/2}},$$

with  $r_\omega = 20h = 2 \text{ kpc}$ , and  $\omega_0$  set to yield the circular rotation speed  $\overline{U}_\phi = 600ht_d^{-1} = 250 \text{ km s}^{-1}$  at  $r = 100h = 10 \text{ kpc}$ . We choose the co-rotation radius to be located at  $r_{\text{cor}} = 100h = 10 \text{ kpc}$ , which corresponds to a pattern speed  $\Omega = 6t_d^{-1} = 25 \text{ km s}^{-1} \text{ kpc}^{-1}$ .

Furthermore, we take  $\alpha_0 = 2.4ht_d^{-1} = 1 \text{ km s}^{-1}$  at all radii. We also take  $\kappa = -0.1h^{-1} = -1 \text{ kpc}^{-1}$  and  $\epsilon_\alpha = 1$ , the latter chosen to be large to maximize the strength of the non-axisymmetric modes. We also adopt  $c_\tau = 1$ .

The value of  $\tau$  is estimated as  $\tau \approx 2l/u \approx 2l^2/3\eta_t \approx 2l^2t_d/3h^2 = t_d/24 = 9.8 \text{ Myr}$ . We also consider solutions for  $\tau \rightarrow 0$ , both for reference and also because  $\tau$  could be much

smaller than our estimate of  $2l/u$  in some galaxies (though it could be much larger in others).

When describing the geometry of the magnetic modes, it is useful to have in mind that magnetic lines have the shape of a spiral with the pitch angle defined in Eq. (30). These spirals have to be distinguished from the spirals along which magnetic field strength is maximum. These can be called magnetic arms or magnetic ridges, and have a different pitch angle (Baryshnikova et al. 1987; Krasheninnikova et al. 1989) (see also Fig. VII.8 of RSS88).

The converged results are shown in Fig. 1 for  $\tau \rightarrow 0$  and  $\tau = t_d/24 = 9.8 \text{ Myr}$ . (Solutions for intermediate values of  $\tau$  are intermediate between the two solutions illustrated.) Panel a shows the potential  $V(r)$  of Eq. (39), along with the ‘energy’  $-(1+\gamma\tau)\gamma t_d/h^2$  of the fastest growing eigenmode. Importantly, due to the non-axisymmetric part of  $\alpha$ , the potential has a local minimum near the co-rotation radius. Therefore, corotating non-axisymmetric modes can be preferentially excited there. The finite value of  $\tau$  makes the potential deeper and thus enhances the growth of non-axisymmetric modes, with a slightly larger growth rate,  $\Gamma \simeq 1.41t_d^{-1} = 5.9 \text{ Gyr}^{-1}$  for  $\tau = t_d/24$  versus  $\Gamma \simeq 1.33t_d^{-1} = 5.6 \text{ Gyr}^{-1}$  for  $\tau \rightarrow 0$ .

Figure 1b shows the ratio

$$\delta \equiv \frac{\overline{B}_\phi - \overline{B}_\phi^{(0)}}{\overline{B}_\phi^{(0)}} = \delta^{(2)}, \quad (42)$$

where the last equality does not hold for numerical solutions, which include higher-order modes, and where,

$$\delta^{(m)} \equiv \frac{\overline{B}_\phi^{(m)}}{\overline{B}_\phi^{(0)}} \cos[m(\phi - \phi_\phi)]. \quad (43)$$

The quantity  $\delta^{(2)}$  is given by the second term in the brackets of Eq. (29). The figures show  $\delta$  at the azimuth  $\phi = \phi_{\text{cor}} = \kappa r_{\text{cor}}/2$  where one of the  $\alpha$ -spiral arms crosses the co-rotation circle. The  $(\phi$ -independent) amplitude (envelope) of this ratio is shown with dashed curves, and  $\alpha(r, \phi_{\text{cor}})$  is represented with a dash-dotted curve for reference. For both finite and vanishing  $\tau$ , the envelope of  $\delta$  exceeds unity near the co-rotation, so that the  $|m| = 2$  modes dominate there. The dominance is stronger when  $\tau$  differs from zero. Importantly, the radius  $r_{\text{max}}$  where the envelope of  $\delta$  is maximum is somewhat larger for the finite  $\tau$ .

The azimuthal phase differences  $\Delta_\phi$  and  $\Delta_r$  between the maximum in, respectively, the azimuthal and radial magnetic field components, and the maximum of  $\alpha$ , are shown in Fig. 1c. For both  $\tau \rightarrow 0$  and  $\tau \neq 0$ , the magnetic spiral arms (whose phase shift is given approximately by  $\Delta_\phi$  since  $\overline{B}_\phi$  dominates over  $\overline{B}_r$ ) cross the  $\alpha$ -spiral near the co-rotation radius, so that each magnetic arm leads the  $\alpha$ -arm in azimuth inside the co-rotation circle and lags it at larger radii. This implies that the magnetic arms are more tightly wound than the material arms.

The  $\tau$  effect produces a phase shift in the magnetic arm such that the part of the arm that leads the corresponding  $\alpha$ -arm is weakened, while the part that lags the  $\alpha$ -arm is enhanced. This can be seen from Fig. 1c. For the  $\tau \rightarrow 0$  case,  $r_{\text{max}}$  (vertical dotted red line) is located slightly inside the circles at which  $\Delta_\phi = 0$  and  $\Delta_r = 0$ . At these circles, the  $\overline{B}_\phi$  and  $\overline{B}_r$  magnetic arms respectively cross the  $\alpha$ -arm. This means that for  $\tau \rightarrow 0$ , the part of the magnetic arm

that leads the  $\alpha$ -arm is stronger than that which lags. (For the more realistic numerical solutions of Paper I, the leading and lagging parts are of equal strength for  $\tau \rightarrow 0$  as can be seen in Fig. 8c of that paper.) On the other hand, for  $\tau = 2l/u$  in our asymptotic solution, we see that  $r_{\text{max}}$  is located to the right of the radii for which  $\Delta_\phi = 0$  and  $\Delta_r = 0$ , which tells us that the lagging part of the magnetic arm is stronger than the leading part. This is consistent with the findings of Paper I. Thus, the lagging, outer part of the magnetic arm is enhanced for finite  $\tau$ , while the leading, inner part is weakened. For example, the phase shift between the magnetic and  $\alpha$ -arms is  $\Delta_\phi(r_{\text{max}}) = 5.2^\circ$  for  $\tau \rightarrow 0$  but  $-3.7^\circ$  for  $\tau = 2l/u$ . However, what we are most interested in is the effect of a finite  $\tau$ , so from now on we will mostly consider the phase *difference* between the vanishing and finite  $\tau$  cases. Crucially, we shall show below in Sect. 4.3 that the general behaviour of the phase difference as a function of  $\Omega\tau$  is nicely reproduced by the asymptotic solution. Thus, defining the difference between the phase shifts obtained for  $\tau = 0$  and finite  $\tau$ ,

$$\Delta(\tau) = \Delta_\phi[\tau, r_{\text{max}}(\tau)] - \Delta_\phi[0, r_{\text{max}}(0)], \quad (44)$$

we have  $\Delta(2l/u) = -9^\circ$  for the asymptotic solution, which is of the same order of magnitude as  $-\Omega\tau = -14^\circ$ . The phase difference  $\Delta(\tau) \simeq -\Omega\tau$ , obtained here and in the models of Paper I, is a natural consequence of the finite response time  $\tau$  of  $\mathcal{E}$  to variations in  $\alpha$ .

In Fig. 1d, we plot the pitch angle of the magnetic field  $p_B$  at the same azimuth  $\phi = \phi_{\text{cor}} = \kappa r_{\text{cor}}/2$ , shown with the opposite sign for presentational convenience. The pitch angle obtained for the axisymmetric disc (dashed lines) is about  $p_B = -(20-30)^\circ$  in the region shown, and increases with radius in agreement with Eq. (37). Its large magnitude is due to the large, constant value of  $\alpha_0$  used in this model, whereas the increase of  $|p_B|$  with  $r$  is due to our choice of a constant disc scale height here, as discussed above. The azimuthal variation in  $\alpha$  leads to a large variation in the pitch angle, with  $-p_B$  being large where  $\alpha$  is large and small where  $\alpha$  is small (compare with the dash-dotted curve for  $\alpha$  in Fig. 1b). The magnitude of the variation depends on  $\epsilon_\alpha$ . This variation is explained by the fact that the  $\alpha\omega$  dynamo mechanism tends to produce  $|\overline{B}_r/\overline{B}_\phi| < 1$  due to the velocity shear; hence,  $|p_B| \propto |\alpha_0/G|^{1/2}$ , Eq. (35). (By contrast, the  $\alpha^2$ -dynamo would generate rather open magnetic spirals with  $|\overline{B}_r/\overline{B}_\phi| \simeq 1$ .)

## 4 COMPARISON WITH NUMERICAL RESULTS

### 4.1 Numerical solutions of Paper I

The semi-analytical solution obtained agrees well, qualitatively, with the numerical solutions of Paper I, which can be seen by comparing Fig. 1 with Fig. 8 of Paper I, which shows the solution of the fiducial non-axisymmetric model of that paper (Model E) in the kinematic regime. This is especially true in the  $\tau \rightarrow 0$  case. The numerical solution, however, does reveal a greater difference between the finite and vanishing  $\tau$  cases than what is seen in the semi-analytical solution. For example, the phase difference for Model E of Paper I during the kinematic stage is  $\Delta(l/u) = -27^\circ \pm 4^\circ$




**Figure 2.** The relative strength  $\delta$  of the non-axisymmetric part of the mean magnetic field in Model E of Paper I, defined as in Eq. (42), as a function of the galactocentric radius near the co-rotation circle at  $\phi = \phi_{\text{cor}}$  (black solid), and the envelope of this dependence at all values of  $\phi$  (black dotted). The individual contributions of the  $|m| = 2$  [red dashed for  $\delta^{(2)}(r_{\text{cor}})$  and red dotted for its envelope] and  $|m| = 4$  [blue dash-dotted for  $\delta^{(4)}(r_{\text{cor}})$  and blue dotted for its envelope] modes are also shown. (a)  $\tau \rightarrow 0$ ; (b)  $\tau = l/u$ .

where the range of  $\Delta$  reflects the finite resolution of the numerical grid. This is comparable to, though somewhat larger in magnitude, than  $-\Omega\tau = -18^\circ$  for that model, while in our semi-analytical solution, presented above,  $\Delta$  was found to be comparable, but somewhat smaller in magnitude, than  $-\Omega\tau$  for the disc model used. This apparent discrepancy suggests that there is an additional physical element missing in the asymptotic solution. In fact, the larger phase shift in the numerical solution can be partly attributed to higher-order

even corotating modes, especially  $m = \pm 4$ . Figure 2a shows the relative strength of the non-axisymmetric magnetic field  $\delta$ , defined in Eq. (42), at the azimuthal angle  $\phi = \phi_{\text{cor}}$ , where the  $\alpha$ -spiral crosses the co-rotation circle, for the case  $\tau \rightarrow 0$  in Model E of Paper I. Fig. 2b shows  $\delta(\phi_{\text{cor}})$  for the case  $\tau = l/u$ . These panels are similar to Fig. 8b of Paper I, with the envelope of  $\delta$  over all  $\phi$  plotted here as a dotted line. The quantities  $\delta^{(2)}$  and  $\delta^{(4)}$  [see Eq. 43], are also plotted. It is clear that the  $|m| = 2$  modes dominate, as assumed in the asymptotic solution. However, it is also clear that the ratio of the amplitude of  $|m| = 4$  to that of  $|m| = 2$  is larger when  $\tau$  is finite. Moreover, for  $\tau = l/u$ , the  $\delta^{(4)}$  envelope peaks at a larger radius than that of  $\delta^{(2)}$ . This suggests that the  $|m| = 4$  modes have the effect of shifting  $r_{\text{max}}$  outward. Since  $\Delta(\tau)$  is sensitive to  $r_{\text{max}}(\tau)$ , this can cause a significant increase in  $\Delta(\tau)$ .

Although one could, in principle, include modes with  $|m| > 2$  in the asymptotic solution presented above, every extra mode would add four new differential equations as well as new terms to the existing equations (16)–(19), greatly increasing the complexity of the model. We take the view that the asymptotic solution presented above does a very good job of reproducing the key qualitative features of the numerical solution, and that further refinements for the sake of numerical accuracy are not necessary at this time.

#### 4.2 Direct comparison of asymptotic and numerical solutions

It is also useful to compare an asymptotic solution such as that presented above with a numerical solution for *the same disc model* (with e.g. a much smaller half-thickness  $h$  than the models of Paper I). An example is shown in Figs. 3 and 4 for  $\tau \rightarrow 0$  and  $\tau = l/u$ , respectively. The model uses the same parameters as the asymptotic solution above, except that  $\epsilon = 0.6$  instead of 1, for both the asymptotic and numerical solutions. Plots are shown at  $t = 10t_d$ , when the relative strength of higher-order *radial* modes has become quite low (though they can still be seen as wiggles at  $r > 11$  kpc). In Figs. 3a (asymptotic) and c (numerical), we compare the quantity  $\delta$  for the two types of solution, while in 3b (asymptotic) and d (numerical), the quantities  $\Delta_r$  and  $\Delta_\phi$  are compared, for the  $\tau \rightarrow 0$  case. For this case of vanishing  $\tau$ , the two types of solution are in fairly good agreement. However, as can be seen by comparing panels a and c, magnetic arms are more pronounced in the numerical solution than in the asymptotic solution. This can be attributed to higher order enslaved modes in the numerical solution; as seen by comparing the overall envelope (dashed) with the envelopes of the  $|m| = 2$  (dotted) and  $|m| = 4$  (dash-dotted) modes in panel c.

For the  $\tau = l/u$  case, shown in Fig. 4, qualitative features of the solution are in reasonable agreement, but the effect of a finite  $\tau$  is more dramatic in the numerical solution than in the asymptotic solution. For instance,  $r_{\text{max}}$  is much larger in the numerical solution, resulting in a larger phase shift. This is seen by comparing panels b and d, where it is evident that the black solid and dashed lines at much larger values of  $|\Delta_\phi|$  and  $|\Delta_r|$  in the numerical solution. As with the solutions from Paper I, enslaved modes with  $|m| > 2$  are stronger in the finite  $\tau$  case than in the  $\tau \rightarrow 0$  case, as can be seen




**Figure 3.** Direct comparison of asymptotic (a-b) and numerical (c-d) solutions for the same disc model, for the case  $\tau \rightarrow 0$ . Panels a and c are similar to panel b of Fig. 1, whilst panels b and d are similar to panel c of that figure. In a and c the  $|m| = 2$  and  $|m| = 4$  envelopes are represented by dotted and dash-dotted curves, respectively. In Panel d, the curves are smoothed over 0.125 kpc.

by comparing the dash-dotted envelopes for the  $|m| = 4$  modes in Figs. 4c and 3c. The assertion, made above, that higher-order enslaved modes are responsible for enhancing the phase shift is therefore strengthened.

The fact that non-axisymmetric modes can be easier generated in a thinner disc, where the difference in the global mode structure is less important because of the stronger dominance of the local dynamo action (due to the shorter magnetic diffusion time across the disc), is well known (Sect. VII.8 in RSS88). However, the importance of the finite relaxation time in this respect was not appreciated earlier. We emphasize that this disc model is used for the sake of illustration only, and is not as realistic as the models of Paper I, where enslaved modes with  $|m| > 2$  are anyway much less important.

#### 4.3 The phase difference caused by a finite dynamo relaxation time

It is also interesting to explore how the phase difference  $\Delta$  varies with the dimensionless quantity  $\Omega\tau$ . Here we retain the disc model (of the asymptotic example or of Model E from Paper I) but vary  $\tau$ . Figure 5 shows the phase difference  $\Delta$  as a function of  $\Omega\tau$  for the asymptotic solution of Sect. 3.3, shown by crosses in the plot, confirming that  $\Delta \approx -C\Omega\tau$  with  $C$  a constant of order unity. It can be seen that the relation is not strictly linear and tends to flatten as  $\Omega\tau$  increases, but is approximately linear for  $\Omega\tau < 0.5$ , which is the region of parameter space that normally applies to disc galaxies. Figure 5 also presents results from the numerical solution of Model E from Paper I, discussed in Sect. 4.1 above, where the galaxy model is more realistic than in the




**Figure 4.** Same as Fig. 3 (but note the change in the plotting range of  $\delta$ ), now for  $\tau = l/u$ .

analytical solution obtained in Sect. 3.3. For Model E, the scaling relation  $\Delta \propto \Omega\tau$  (for small  $\Omega\tau$ ) quickly establishes itself in the kinematic regime, as soon as the leading eigenfunction becomes dominant. Remarkably, it applies to both kinematic dynamo solutions and to the steady state, with more or less the same proportionality constant. The magnitude of the proportionality constant  $C \approx 1.7$  (represented by the solid line in the figure) is still of order unity, but is significantly larger than  $C \approx 0.65$  (dashed line) found in the asymptotic solution. Although this difference may be partly attributable to the different disc models used, the larger value of  $C$  in the numerical model is also partly due to higher-order even modes, as discussed in Sect. 4.1 above.

In summary, asymptotic solutions obtained here clarify numerical results of Paper I and help us to isolate their generic features. Figure 1 can be compared with Fig. 8 of Paper I, which shows similar kinematic results, confirming the presence of a strong  $|m| = 2$  magnetic component near  $r = r_{\text{cor}}$ . Around this radius we see clearly from the asymptotic analysis that the effective potential  $V(r)$  has a new minimum, allowing new bound states (or growing modes) of  $q(r)$  near co-rotation. Both asymptotic and numerical solutions reveal stationary even modes in the frame corotating with the spiral pattern of the  $\alpha$ -coefficient. Both have magnetic spiral arms more tightly wound than the  $\alpha$ -arms, with the two types of arm crossing near the co-rotation circle. Moreover, both numerical and asymptotic results, the latter, e.g., in Eq. (34), also agree in that the response of the mean magnetic field to the  $\alpha$ -spiral pattern is stronger when  $\tau$  is finite. More importantly, they agree in that magnetic arms undergo a negative phase shift of order  $-\Omega\tau$  such that the part of the magnetic arm which trails the  $\alpha$ -arm is enhanced when  $\tau$  is finite. Enslaved modes with  $|m| > 2$ , though weak, can substantially enhance the phase shift because they are located further out in radius than the  $|m| = 2$  mode, where magnetic arms lag  $\alpha$ -arms.




**Figure 5.** The phase difference  $\Delta$  between the solutions with  $\tau \neq 0$  and  $\tau = 0$  as a function of  $\Omega\tau$ . Results for the parameter values used in the illustrative example are shown with blue crosses; those from Model E of Paper I are shown during the kinematic regime (at  $t = 3t_{d,0}$  after the simulation has begun) and in the steady state.

## 5 CONCLUSIONS

We have extended non-axisymmetric mean-field dynamo theory in two important ways:

- Obtained semi-analytically a solution for the axisymmetric and strongest enslaved non-axisymmetric modes ( $m = 0$  and  $m = \pm 2$  for a two-armed spiral). Previously, the only analytical treatment that existed was for the subdominant non-enslaved modes ( $m = \pm 1$ );
- Included the effects of the finite relaxation time of the mean electromotive force (the  $\tau$  effect) in our solutions.

The asymptotic solutions employ the WKB approximation applicable to tightly wound magnetic spirals; to obtain explicit expressions for the growth rate of the mean magnetic field and its radial and azimuthal distribution we use the no- $z$  approximation (known to be quite accurate for thin discs) to solve the local dynamo equations.

We find overall good qualitative agreement with the numerical results of Paper I for a global, rigidly rotating spiral pattern (which is the type of spiral forcing most amenable to analytical treatment) to grow around the co-rotation radius. In particular, the asymptotic solution obtained here agrees with the numerical solution of Paper I in the following key features.

(i) Strong magnetic modes with the azimuthal wave numbers  $m = \pm n$  are supported by the dynamo action, which corotate with the  $\alpha$ -spiral (here  $n$  is the number of  $\alpha$ -spiral arms). Non-axisymmetric modes are confined to a radial range of a few kiloparsecs around the co-rotation radius and are comparable in strength with the  $m = 0$  mode.

(ii) Magnetic arms (understood as ridges of the mean magnetic field strength that have a spiral shape) cross the  $\alpha$ -spiral near the co-rotation radius and are more tightly wound than the  $\alpha$ -arms.

(iii) In the model considered here, the only effect of the galactic spiral pattern is the enhancement of the dynamo  $\alpha$ -coefficient by the spiral pattern. As a result, the magnetic lines of the mean field are less tightly wound (larger magnitude of the pitch angle  $p_B$ ) within the  $\alpha$ -spirals than between them.

(iv) The amplitude of the non-axisymmetric modes increases with the magnitude of the relaxation time  $\tau$ .

(v) Another effect of the finite magnitude of  $\tau$  is that the part of the magnetic arm that lags behind the  $\alpha$ -arm with respect to the galactic rotation (i.e., outside the co-rotation radius) becomes stronger than the part that leads the  $\alpha$ -arm (inside the co-rotation radius). This leads to the overall appearance of magnetic arms that are phase-shifted from the  $\alpha$ -arms (material arms) in the direction opposite to the galactic rotation.

(vi) For  $\Omega\tau \lesssim 0.5$ , the phase shift produced by a finite value of  $\tau$  (relative to that for  $\tau \rightarrow 0$ ), is directly proportional to  $-\Omega\tau$ , with a proportionality constant of order unity.

## 6 DISCUSSION FOR PAPERS I AND II

Results obtained in Chamandy et al. (2013) (Paper I) and here (Paper II) for a rigidly rotating  $\alpha$ -spiral demonstrate the ability of the mean-field dynamo mechanism to produce magnetic arms that do not overlap with the material arms in the regions where the former are best pronounced, which is outside the co-rotation circle in our model. Due to a finite magnitude of the relaxation time of the mean electromotive force,  $\tau$ , the lagging part of the magnetic arm is better pronounced than that ahead of the material arm, which can make it difficult to detect their intersection in the observations. As a result of advection of the mean magnetic field by the differentially rotating gas, magnetic arms are, in these models, more tightly wound than the  $\alpha$ -spiral which drives them, and are localized to within a few kiloparsecs of the co-rotation circle. The phase shift between the magnetic and  $\alpha$ -arms in our model vanishes at or near the co-rotation radius and then increases, first rapidly and then at a lower rate, up to a value as large as  $60^\circ$  at a distance of a few kpc from the co-rotation circle.

The morphology of magnetic arms in spiral galaxies and their position relative to the material arms is rather diverse. One extreme is the galaxy NGC 6946 where magnetic arms appear to have the same pitch angle as the optical arms, and the two patterns do not intersect. The azimuthal phase shifts between the individual optical arms and their magnetic counterparts identified by Frick et al. (2000) are  $-26^\circ \pm 12^\circ$ ,  $-36^\circ \pm 11^\circ$  and  $-45^\circ \pm 17^\circ$  at  $r \approx 4$  kpc and, for an outer arm,  $-68^\circ \pm 17^\circ$  at  $r \approx 8$  kpc, where negative phase shift corresponds to a position behind a gaseous arm (all distances have been reduced to a distance of 5.5 Mpc to NGC 6946 – Kennicutt et al. 2003). According to the H $\alpha$  observations and analysis of Fathi et al. (2007), the co-rotation radius of the outer spiral pattern in this galaxy is at about 8.3 kpc, whereas the interlaced magnetic and optical spiral arms occur at  $1 \lesssim r \lesssim 9$  kpc. The dynamo action at  $r \gtrsim 9$  kpc may be too weak to make the magnetic arms observable there, but there are no indications of the magnetic and optical arms crossing at or near the co-rotation radius. Nevertheless, the right magnitude of the azimuthal phase shifts obtained in our models ( $15$ – $40^\circ$ ) is arguably encouraging.

Another possibility is that the spiral structure in NGC 6946 is more complex than that of a single, rigidly rotating pattern. For instance, the arms may be winding

up to some extent, and thus transient (e.g. Dobbs et al. 2010; Sellwood 2011; Quillen et al. 2011; Wada et al. 2011; Grand et al. 2012). The model of Comparetta & Quillen (2012) invokes multiple rigidly rotating patterns, which interfere to produce spiral features that wind up. Magnetic arms which are present over several kiloparsecs in radius, and which have a large negative phase shift from the  $\alpha$ -arms that varies only weakly with radius, like in NGC 6946, are indeed found in our ‘winding-up’ spiral model of Paper I.

On the other hand, the mutual arrangement of magnetic and material arms is more complicated in M51 (Patrikeev et al. 2006; Fletcher et al. 2011) where the two patterns overlap in some regions but are systematically offset by about 0.5–0.6 kpc elsewhere (adopting 7.6 Mpc for the distance to M51 – Ciardullo et al. 2002). This linear displacement corresponds to the angular phase difference of 5–10° at  $r = 3$ –6 kpc where such an offset is best pronounced. The two arms visible in polarized emission at  $\lambda 6.2$  cm intersect the material arms observed in other tracers at  $r = 5$ –6 kpc. Elmegreen et al. (1989) suggest that M51 has two spiral arm systems, with the co-rotation radii at  $r = 4.8$  kpc and  $r = 12$  kpc. García-Burillo et al. (1993) determine the co-rotation radius to be 5.8 kpc from CO observations and modelling of M51. Then the region at  $r = 3$ –6 kpc is (mostly) inside the co-rotation radius and the magnetic arms in the inner galaxy are displaced downstream of the material arms, as expected in our model. The two magnetic arms are positioned differently with respect to the material arms at larger galactocentric distances. Outside the co-rotation circle, magnetic Arm 1 (on the east of the galactic centre) is systematically lagging the material arm, in accordance with our model, but the magnetic Arm 2 overlaps the material arm.

Our model appears to be better applicable to this galaxy, but the diverse mutual arrangement of the magnetic and material arms in M51 strongly suggests that more than one physical effect is involved in its genesis. Dobbs et al. (2010) suggest that the spiral arm morphology in M51 evolves rapidly due to the interaction with its satellite galaxy. Magnetic arms driven by evolving material arms are explored in Paper I. It also appears that M51 may be a good candidate for the ‘ghost’ magnetic arms, where the magnetic arms trace the spiral arms as they were in the past up to a few hundred Myr earlier. The simulations of Dobbs et al. (2010) show that the material spiral arms are being wound up by the galactic differential rotation, so the model with transient material spirals of Paper I may be more relevant to M51 than the model with a rigidly rotating, stationary spiral pattern explored here.

Still another morphology of magnetic and gaseous arms is observed in the nearby barred galaxy M83. Beck et al. (in preparation) applied the same approach involving wavelet transforms as Frick et al. (2000) to determine the positions of the spiral arms as seen in optical light, dust, H $\alpha$ , CO, as well as the total and polarized intensities at  $\lambda\lambda 6$  and 13 cm. The morphology of M83 is dominated by a well-pronounced two-armed spiral pattern in each tracer, with some substructure within the arms. The magnetic and material arms in this galaxy are clearly separated but intersect at the galactocentric radius 6.4 kpc (given the distance to M83 is 4.5 Mpc – Thim et al. 2003). The co-rotation radius of M83, determined by Hirota et al. (2009) as 2.4 arcmin, or 6.2 kpc, is practically equal to the intersec-

tion radius given the uncertainties involved. This galaxy is a good candidate for the formation of magnetic arms by the mechanism suggested here.

An important assumption of all or most of the available models of magnetic arms is that the  $\alpha$ -effect is stronger within, or at least correlated with, the material spiral arms. We note that the effects of the spiral arms on the scale height of the interstellar gas, its turbulent scale and velocity, local velocity shear, and other parameters that control the intensity of the dynamo action are far from being certain, either observationally or theoretically (see Shukurov 1998; Shukurov et al. 2004, and references therein). Shukurov & Sokoloff (1998) argue that the magnitude of the  $\alpha$  effect within the material arms can be four times smaller than between the arms, whereas the turbulent diffusivity is plausible to be only weakly modulated by the spiral pattern. As a result, the local dynamo number within the material arms can be four times smaller than between them, and the dynamo action can, in fact, be more vigorous between the material arms. Direct numerical simulations by Elstner & Gressel (2012) find that both the  $\alpha$  effect and turbulent diffusivity increase with star formation rate  $\sigma$ , which itself will be larger within the arms, but this leads to an overall dynamo number that scales inversely with  $\sigma$ . However, even if the dynamo number is larger in the interarm regions, this does not necessarily imply that the steady-state mean magnetic field there should be stronger than within the arms, since the gas density is lower in between the arms than within them. Significant effort in observations, theory of interstellar magnetohydrodynamics and dynamo theory are still required to establish a clear understanding of the various mechanisms that produce the diverse morphology of magnetic arms in spiral galaxies. Our work here and in Paper I provides the beginnings of such a study.

## APPENDIX A: THE LOCAL EIGENFUNCTIONS

With approximations described in Section 3.2.1, Eqs. (18) and (19) yield, after some algebra,

$$b_{\pm 2} = b_0 \frac{\alpha_{\pm 2}}{\alpha_0} |D_0| X_{\pm 2}, \quad a_{\pm 2} = -Y_{\pm 2} b_{\pm 2}. \quad (\text{A1})$$

Here

$$X_{\pm 2}(r) = \frac{(A \mp iB)}{A^2 + B^2} = \frac{e^{\mp i\beta}}{\sqrt{A^2 + B^2}} = X_{\mp 2}^*, \quad (\text{A2})$$

and

$$Y_{\pm 2}(r) = \tilde{A} \mp i\tilde{B} = \sqrt{\tilde{A}^2 + \tilde{B}^2} e^{\mp i\tilde{\beta}} = Y_{\mp 2}^*, \quad (\text{A3})$$

with the dynamo number defined as

$$D_0 = \frac{\alpha_0 G h^3}{\eta_t^2} = \frac{\alpha_0 G t_d^2}{h} < 0, \quad (\text{A4})$$

where  $t_d = h^2/\eta_t$  is the vertical turbulent diffusion time scale and

$$\cos \beta = \frac{A}{\sqrt{A^2 + B^2}}, \quad \sin \beta = \frac{B}{\sqrt{A^2 + B^2}}, \quad (\text{A5})$$

with

$$A = D_0 + \frac{\pi}{2c_\tau} \left\{ t_d^2 (1 + \Gamma\tau)(\Gamma^2 - 4\tilde{\omega}^2) + 8\Omega\Gamma\tilde{\omega}\tau t_d^2 \right. \\ + [(1 + \Gamma\tau)^2 + (2\Omega\tau)^2]^{-1} \\ \times \left[ c_\tau^2 (1 + \Gamma\tau) \left( \frac{1}{4}\pi^2 + \theta_a'^2 h^2 \right) \left( \frac{1}{4}\pi^2 + \theta_b'^2 h^2 \right) \right. \\ - 2c_\tau^2 \Omega\tau \left[ \left( \frac{1}{4}\pi^2 + \theta_a'^2 h^2 \right) \theta_b'' h^2 + \left( \frac{1}{4}\pi^2 + \theta_b'^2 h^2 \right) \theta_a'' h^2 \right] \\ \left. - c_\tau^2 (1 + \Gamma\tau) \theta_a'' \theta_b'' h^4 \right] \\ \left. + c_\tau \Gamma t_d \left( \frac{1}{2}\pi^2 + \theta_a'^2 h^2 + \theta_b'^2 h^2 \right) - 2c_\tau \tilde{\omega} t_d h^2 (\theta_a'' + \theta_b'') \right\}, \quad (\text{A6})$$

$$B = \frac{\pi}{2c_\tau} \left\{ -2\Omega\tau t_d^2 (\Gamma^2 - 4\tilde{\omega}^2) + 4(1 + \Gamma\tau)\Gamma\tilde{\omega} t_d^2 \right. \\ + [(1 + \Gamma\tau)^2 + (2\Omega\tau)^2]^{-1} \\ \times \left[ 2c_\tau^2 \Omega\tau \left( \frac{1}{4}\pi^2 + \theta_a'^2 h^2 \right) \left( \frac{1}{4}\pi^2 + \theta_b'^2 h^2 \right) \right. \\ + c_\tau^2 (1 + \Gamma\tau) \left[ \left( \frac{1}{4}\pi^2 + \theta_a'^2 h^2 \right) \theta_b'' h^2 + \left( \frac{1}{4}\pi^2 + \theta_b'^2 h^2 \right) \theta_a'' h^2 \right] \\ \left. - 2c_\tau^2 \Omega\tau \theta_a'' \theta_b'' h^4 \right] \\ \left. + 2c_\tau \tilde{\omega} t_d \left( \frac{1}{2}\pi^2 + \theta_a'^2 h^2 + \theta_b'^2 h^2 \right) + c_\tau \Gamma t_d h^2 (\theta_a'' + \theta_b'') \right\}. \quad (\text{A7})$$

Further,

$$\cos \tilde{\beta} = \frac{\tilde{A}}{\sqrt{\tilde{A}^2 + \tilde{B}^2}}, \quad \sin \tilde{\beta} = \frac{\tilde{B}}{\sqrt{\tilde{A}^2 + \tilde{B}^2}}, \quad (\text{A8})$$

with

$$\tilde{A} = -\frac{\Gamma}{G} - c_\tau \frac{(1 + \Gamma\tau) \left( \frac{1}{4}\pi^2 + \theta_b'^2 h^2 \right) - 2\Omega\tau \theta_b'' h^2}{[(1 + \Gamma\tau)^2 + (2\Omega\tau)^2] G t_d}, \quad (\text{A9})$$

$$\tilde{B} = \frac{2\tilde{\omega}}{G} + c_\tau \frac{2\Omega\tau \left( \frac{1}{4}\pi^2 + \theta_b'^2 h^2 \right) + (1 + \Gamma\tau) \theta_b'' h^2}{[(1 + \Gamma\tau)^2 + (2\Omega\tau)^2] G t_d}. \quad (\text{A10})$$

Substituting (15), (A2) and (A3) into Eq. (A1), we arrive at

$$a_{\pm 2} = -\frac{|D_0| b_0 \epsilon_\alpha}{2} \sqrt{\frac{\tilde{A}^2 + \tilde{B}^2}{\tilde{A}^2 + \tilde{B}^2}} \exp \left[ \mp i(\kappa r + \beta + \tilde{\beta}) \right], \quad (\text{A11})$$

$$b_{\pm 2} = \frac{|D_0| b_0 \epsilon_\alpha}{2} \frac{1}{\sqrt{\tilde{A}^2 + \tilde{B}^2}} \exp \left[ \mp i(\kappa r + \beta) \right]. \quad (\text{A12})$$

## REFERENCES

- Baryshnikova Y., Shukurov A., Ruzmaikin A., Sokoloff D. D., 1987, *A&A*, 177, 27
- Beck R., 2012, *SSRv*, 166, 215
- Beck R., Brandenburg A., Moss D., Shukurov A., Sokoloff D., 1996, *ARA&A*, 34, 155
- Beck R., Ehle M., Frick P., Patrikeyev I., Shukurov A., Sokoloff D. D., in preparation, in preparation
- Blackman E. G., Field G. B., 2002, *PRL*, 89, 265007
- Brandenburg A., Subramanian K., 2005a, *PhR*, 417, 1
- , 2005b, *A&A*, 439, 835
- , 2007, *Astron. Nachr.*, 328, 507
- Chamandy L., Subramanian K., Shukurov A., 2013, *MNRAS*, 428, 3569
- Ciardullo R., Feldmeier J. J., Jacoby G. H., Kuzio de Naray R., Laychak M. B., Durrell P. R., 2002, *ApJ*, 577, 31
- Comparetta J., Quillen A. C., 2012, *ArXiv e-prints*
- Dobbs C. L., Theis C., Pringle J. E., Bate M. R., 2010, *MNRAS*, 403, 625
- Elmegreen B. G., Seiden P. E., Elmegreen D. M., 1989, *ApJ*, 343, 602
- Elstner D., Gressel O., 2012, *ArXiv e-prints*
- Eyink G. L., 2012, *Turbulence Theory*, Course Notes, John Hopkins Univ., <http://www.ams.jhu.edu/~eyink/Turbulence/>
- Fathi K., Toonen S., Falcón-Barroso J., Beckman J. E., Hernandez O., Daigle O., Carignan C., de Zeeuw T., 2007, *ApJ*, 667, L137
- Fletcher A., 2010, in *Astronomical Society of the Pacific Conference Series*, Vol. 438, *Astron. Soc. Pacific, Conference Ser.*, Kothes R., Landecker T. L., Willis A. G., eds., p. 197
- Fletcher A., Beck R., Shukurov A., Berkhuijsen E. M., Horellou C., 2011, *MNRAS*, 412, 2396
- Frick P., Beck R., Shukurov A., Sokoloff D., Ehle M., Kamphuis J., 2000, *MNRAS*, 318, 925
- García-Burillo S., Combes F., Gerin M., 1993, *A&A*, 274, 148
- Gent A. F., Shukurov A., Fletcher A., Sarson G. R., Mantere M. J., 2012, *MNRAS*
- Germano M., 1992, *JFM*, 238, 325
- Grand R. J. J., Kawata D., Cropper M., 2012, *MNRAS*, 421, 1529
- Hirota A., Kuno N., Sato N., Nakanishi H., Tosaki T., Matsui H., Habe A., Sorai K., 2009, *PASJ*, 61, 441
- Hubbard A., Brandenburg A., 2009, *ApJ*, 706, 712
- Kennicutt Jr. R. C., Armus L., Bendo G., Calzetti D., Dale D. A., Draine B. T., Engelbracht C. W., Gordon K. D., et al., 2003, *PASP*, 115, 928
- Krasheninnikova Y., Shukurov A., Ruzmaikin A., Sokoloff D., 1989, *A&A*, 213, 19
- Mestel L., Subramanian K., 1991, *MNRAS*, 248, 677
- Moss D., 1995, *MNRAS*, 275, 191
- , 1998, *MNRAS*, 297, 860
- Patrikeyev I., Fletcher A., Stepanov R., Beck R., Berkhuijsen E. M., Frick P., Horellou C., 2006, *A&A*, 458, 441
- Phillips A., 2001, *Geophys. Astrophys. Fluid Dyn.*, 94, 135
- Quillen A. C., Dougherty J., Bagley M. B., Minchev I., Comparetta J., 2011, *MNRAS*, 417, 762
- Rheinhardt M., Brandenburg A., 2012, *Astron. Nachr.*, 333, 71
- Rogachevskii I., Kleeorin N., 2000, *PRE*, 61, 5202
- Rohde R., Beck R., Elstner D., 1999, *A&A*, 350, 423
- Roškar R., Debattista V. P., Quinn T. R., Wadsley J., 2012, *MNRAS*, 426, 2089
- Ruzmaikin A. A., Shukurov A. M., Sokoloff D. D., 1988, *Magnetic fields of galaxies*. Kluwer, Dordrecht
- Sellwood J. A., 2011, *MNRAS*, 410, 1637
- Shukurov A., 1998, *MNRAS*, 299, L21
- , 2005, in *Lecture Notes in Physics*, Berlin Springer Verlag, Vol. 664, *Cosmic Magnetic Fields*, Wiełebinski R., Beck R., eds., p. 113
- Shukurov A., Sarson G. R., Nordlund Å., Gudiksen B., Brandenburg A., 2004, *Ap&SS*, 289, 319

- Shukurov A., Sokoloff D., 1998, *Stud. Geophys. Geod.*, 42, 391  
Subramanian K., Mestel L., 1993, *MNRAS*, 265, 649  
Sur S., Shukurov A., Subramanian K., 2007, *MNRAS*, 377, 874  
Thim F., Tammann G. A., Saha A., Dolphin A., Sandage A., Tolstoy E., Labhardt L., 2003, *ApJ*, 590, 256  
Wada K., Baba J., Saitoh T. R., 2011, *ApJ*, 735, 1